

Solution generating with perfect fluids

David Garfinkle

Department of Physics, Oakland University, Rochester, MI 48309

E.N. Glass

Department of Physics, University of Michigan, Ann Arbor, MI 48109

and Physics Department, University of Windsor, Windsor, Ontario, N9B 3P4 (permanent address)

J.P. Krisch

Department of Physics, University of Michigan, Ann Arbor, MI 48109

Abstract

We apply a technique, due to Stephani, for generating solutions of the Einstein-perfect-fluid equations. This technique is similar to the vacuum solution generating techniques of Ehlers, Harrison, Geroch and others. We start with a “seed” solution of the Einstein-perfect-fluid equations with a Killing vector. The seed solution must either have (i) a spacelike Killing vector and equation of state $P = \rho$ or (ii) a timelike Killing vector and equation of state $\rho + 3P = 0$. The new solution generated by this technique then has the same Killing vector and the same equation of state. We choose several simple seed solutions with these equations of state and where the Killing vector has no twist. The new solutions are twisting versions of the seed solutions.

I. INTRODUCTION

In solving Einstein's equation it is usual to treat spacetimes that have one or more Killing vectors. The existence of a Killing vector reduces Einstein's equation from a 4 dimensional set of equations to a 3 dimensional set. Instead of the spacetime metric one considers an equivalent set of variables: the norm λ and twist potential ω of the Killing vector and a metric γ_{ab} on the 3 dimensional manifold of trajectories of the Killing field. The vacuum Einstein equations then become a set of equations for the quantities $(\lambda, \omega, \gamma_{ab})$. Remarkably, as shown by Ehlers [1] , Harrison [2] , Geroch [3] and others [4] these equations have a symmetry that allows us to generate new solutions: given $(\lambda, \omega, \gamma_{ab})$ solving the equations then $(\tilde{\lambda}, \tilde{\omega}, \gamma_{ab})$ is also a solution where $(\tilde{\lambda}, \tilde{\omega})$ is a simple algebraic function of (λ, ω) . These results have been generalized to the Einstein-Maxwell equations [2,5,6] and have been applied to find many new solutions. (The general invariance transformation was first published in [4] (in the vacuum case with two Killing vectors) and in [6] (in the Einstein-Maxwell case with one Killing vector). Ehlers [1] found a special case of the general vacuum transformation while Harrison [2] found a special case of the general Einstein-Maxwell transformation.)

The vacuum solution generating technique was generalized to the case of perfect fluids by Stephani [7]. This technique is very similar to that for the vacuum Einstein equation. However, it turns out that not all equations of state are suitable for this sort of solution generating. Only two equations of state are compatible with this technique: $P = \rho$ for a spacelike Killing field and $\rho + 3P = 0$ for a timelike Killing field.

In section 2 we define our notation. Sections 3 and 4 contain the new solutions that we have generated using this technique with section 3 treating the $P = \rho$ case and section 4 treating the $\rho + 3P = 0$ case. Section 5 contains a brief discussion of our results.

II. FORMALISM

Our notation is essentially that of refs. [3,7]. We use signature $(-, +, +, +)$ and units where $c = G = 1$. Let (M, g_{ab}) be a solution of the Einstein-perfect fluid equations with energy density ρ and pressure P . Assume that g_{ab} has a Killing vector ξ^a . Define the norm λ and twist ω_a of ξ^a by $\lambda := \xi^a \xi_a$ and

$$\omega_a := \epsilon_{abcd} \xi^b \nabla^c \xi^d \quad . \quad (2.1)$$

We assume that either (i) the Killing vector is orthogonal to the fluid four-velocity and $P = \rho$ or (ii) the Killing vector is parallel to the fluid four-velocity and $\rho + 3P = 0$. From this assumption it follows that $R_{ab} \xi^b = 0$. It then follows that there is a scalar ω such that $\omega_a = \nabla_a \omega$ and that there are forms α_a and β_a satisfying

$$\nabla_{[a} \alpha_{b]} = \frac{1}{2} \epsilon_{abcd} \nabla^c \xi^d \quad , \quad (2.2)$$

$$\xi^a \alpha_a = \omega \quad , \quad (2.3)$$

$$\nabla_{[a} \beta_{b]} = 2 \lambda \nabla_a \xi_b + \omega \epsilon_{abcd} \nabla^c \xi^d \quad , \quad (2.4)$$

$$\xi^a \beta_a = \omega^2 + \lambda^2 - 1 \quad . \quad (2.5)$$

For any constant θ define $\tilde{\lambda}$ and η_a by

$$\lambda / \tilde{\lambda} := \cos^2 \theta + (\omega^2 + \lambda^2) \sin^2 \theta - 2 \omega \sin \theta \cos \theta \quad , \quad (2.6)$$

$$\eta_a := \tilde{\lambda}^{-1} \xi_a + 2 \cos \theta \sin \theta \alpha_a - \sin^2 \theta \beta_a \quad . \quad (2.7)$$

Then the new metric is given by

$$\tilde{g}_{ab} = \frac{\lambda}{\tilde{\lambda}} \left(g_{ab} - \lambda^{-1} \xi_a \xi_b \right) + \tilde{\lambda} \eta_a \eta_b \quad . \quad (2.8)$$

This new metric is also a solution of the Einstein-perfect fluid equations with the same equation of state and the same Killing vector.

One of the uses of solution generating techniques is that they can produce twisting solutions from seed solutions that are twist free. We now consider how the process for

finding the new metric simplifies in the case where the seed metric is twist free. In this case $\omega = 0$ and $\nabla_a \xi_b = \lambda^{-1} \xi_{[b} \nabla_{a]} \lambda$. Define $F := \cos^2 \theta + \lambda^2 \sin^2 \theta$. Then $\tilde{\lambda} = \lambda/F$. The equation for β_a can be solved as $\beta_a = (\lambda - \lambda^{-1}) \xi_a$. The equation for α_a simplifies to

$$\nabla_{[a} \alpha_{b]} = \frac{1}{2\lambda} \epsilon_{abcd} \xi^d \nabla^c \lambda \quad (2.9)$$

and $\xi^a \alpha_a = 0$. The one-form η_a is then given by

$$\eta_a = \lambda^{-1} \xi_a + 2 \cos \theta \sin \theta \alpha_a \quad . \quad (2.10)$$

The new metric is then given by

$$\tilde{g}_{ab} = F \left(g_{ab} - \lambda^{-1} \xi_a \xi_b \right) + \frac{\lambda}{F} \eta_a \eta_b \quad . \quad (2.11)$$

From now on we will assume that the seed solution is twist-free.

III. $P = \rho$ FLUIDS

A. Fluid variables

In describing the new solutions we will concentrate on two properties of interest: singularities and fluid flow. The solution generating procedure often results in singularities on the axis. For each of our solutions we will consider whether such a singularity exists. A perfect fluid is described by the energy density ρ , pressure P and fluid four velocity u^a . The covariant derivative of the fluid four velocity is decomposed [8] as

$$\nabla_b u_a = -a_a u_b + \omega_{ab} + \sigma_{ab} + \Theta (g_{ab} + u_a u_b) / 3 \quad . \quad (3.1)$$

Here $a_a := u^b \nabla_b u_a$ is called the acceleration. The tensors σ_{ab} (called the shear) and ω_{ab} (called the vorticity) are orthogonal to u^a with ω_{ab} antisymmetric and σ_{ab} symmetric and trace-free. The scalar Θ is called the expansion. Under solution generating with the $P = \rho$ equation of state the fluid parameters transform as follows:

$$\tilde{\rho} = \rho / F \quad , \quad (3.2)$$

$$\tilde{u}_a = \sqrt{F} u_a \quad , \quad (3.3)$$

$$\tilde{\Theta} = \frac{1}{\sqrt{F}} \left(\Theta + F^{-1} \lambda \sin^2 \theta u^a \nabla_a \lambda \right) \quad , \quad (3.4)$$

$$\tilde{a}_a = a_a + F^{-1} \lambda \sin^2 \theta \left(\nabla_a \lambda + u_a u^b \nabla_b \lambda \right) \quad , \quad (3.5)$$

$$\tilde{\omega}_{ab} = \sqrt{F} \omega_{ab} \quad . \quad (3.6)$$

There does not seem to be a simple transformation rule for σ_{ab} .

A fluid with the equation of state $P = \rho$ is called “stiff matter.” In such a fluid the adiabatic sound speed is equal to the speed of light. If the fluid is irrotational then the field equations are the same as those of a minimally coupled scalar field with timelike gradient.

[9] There is a technique for generating stiff matter solutions from vacuum solutions. [10]

B. Robertson-Walker seed metric

Consider the seed metric

$$ds^2 = - dt^2 + t^{2/3} \left[dz^2 + dr^2 + r^2 d\phi^2 \right] \quad . \quad (3.7)$$

This is a flat Friedmann-Robertson-Walker metric with $P = \rho$. The fluid four velocity is $u_a = -\nabla_a t$ and the energy density is $\rho = 1/(24\pi t^2)$. The shear, vorticity and acceleration all vanish. The expansion is $\Theta = 1/t$. The metric is type O. This metric has more than one Killing vector. We first consider the case $\xi^a = (\partial/\partial z)^a$. Then we have $\lambda = t^{2/3}$ and $\omega = 0$. It then follows that $F = \cos^2 \theta + t^{4/3} \sin^2 \theta$. The equation for α_a is then

$$\nabla_{[a} \alpha_{b]} = -\frac{2}{3} r \nabla_{[a} r \nabla_{b]} \phi \quad (3.8)$$

(as well as $\xi^a \alpha_a = 0$). A solution is

$$\alpha_a = -\frac{1}{3} r^2 \nabla_a \phi \quad . \quad (3.9)$$

The one-form η_a is then given by

$$\eta_a = \nabla_a z - \frac{2}{3} \cos \theta \sin \theta r^2 \nabla_a \phi \quad . \quad (3.10)$$

The new metric $\tilde{g}_{ab}$ is then given by

$$d\tilde{s}^2 = F \left[-dt^2 + t^{2/3} (dr^2 + r^2 d\phi^2) \right] + \frac{t^{2/3}}{F} \left(dz - \frac{2}{3} \cos \theta \sin \theta r^2 d\phi \right)^2 . \quad (3.11)$$

This new metric has no singularities on the axis. The acceleration and vorticity vanish. The expansion is

$$\tilde{\Theta} = \frac{1}{\sqrt{F}} \left(t^{-1} + \frac{2}{3} F^{-1} t^{1/3} \sin^2 \theta \right) . \quad (3.12)$$

The shear has components in the rr , $\phi\phi$, zz and $z\phi$ directions. Its square is

$$\tilde{\sigma}^{ab} \tilde{\sigma}_{ab} = \left(\frac{32}{27} \right) \frac{t^{2/3} \sin^4 \theta}{F^3} . \quad (3.13)$$

The metric is type I.

We now use the same seed metric but with the Killing vector $\xi^a = (\partial/\partial\phi)^a$. Then we have $\lambda = t^{2/3} r^2$ and $\omega = 0$. It then follows that $F = \cos^2 \theta + t^{4/3} r^4 \sin^2 \theta$. The equation for α_a is then

$$\nabla_{[a} \alpha_{b]} = \frac{2}{3} r \nabla_{[a} r \nabla_{b]} z + 2 t^{1/3} \nabla_{[a} t \nabla_{b]} z \quad (3.14)$$

(as well as $\xi^a \alpha_a = 0$). A solution is

$$\alpha_a = \left(\frac{1}{3} r^2 + \frac{3}{2} t^{4/3} \right) \nabla_a z . \quad (3.15)$$

The one-form η_a is then given by

$$\eta_a = \nabla_a \phi + \left(\frac{2}{3} r^2 + 3 t^{4/3} \right) \cos \theta \sin \theta \nabla_a z . \quad (3.16)$$

The new metric $\tilde{g}_{ab}$ is then given by

$$\begin{aligned} d\tilde{s}^2 = F \left[-dt^2 + t^{2/3} (dz^2 + dr^2) \right] \\ + \frac{t^{2/3} r^2}{F} \left(d\phi + \left[\frac{2}{3} r^2 + 3 t^{4/3} \right] \cos \theta \sin \theta dz \right)^2 . \end{aligned} \quad (3.17)$$

This metric has no singularities on the axis. (There is a conical singularity if the range of ϕ is 2π . However, this singularity can be removed by choosing the appropriate range of ϕ).

The vorticity vanishes. The expansion and acceleration are given by

$$\tilde{\Theta} = \frac{1}{\sqrt{F}} \left(t^{-1} + \frac{2}{3} F^{-1} t^{1/3} r^4 \sin^2 \theta \right) , \quad (3.18)$$

$$\tilde{a}_a = 2F^{-1}t^{4/3}r^3\sin^2\theta\nabla_a r \quad . \quad (3.19)$$

The shear has components in the zz , rr , $\phi\phi$ and $z\phi$ directions. Its square is

$$\tilde{\sigma}^{ab}\tilde{\sigma}_{ab} = \left(\frac{8}{27}\right) \frac{r^2 t^{2/3} \sin^2\theta}{F^3} \left[27 \cos^2\theta + 4r^6 \sin^2\theta\right] \quad . \quad (3.20)$$

The metric is type I.

C. Tabensky-Taub seed metric

We now consider a different seed metric: The Tabensky-Taub metric [9]

$$ds^2 = -V dt^2 + V dz^2 + z(dx^2 + dy^2) \quad (3.21)$$

where

$$V = z^{-1/2} \exp\left(\frac{1}{2}a^2 z^2\right) \quad . \quad (3.22)$$

This is a solution of the Einstein-perfect-fluid equations with

$$P = \rho = \frac{a^2}{16\pi V} \quad . \quad (3.23)$$

The fluid four-velocity is $u_a = -\sqrt{V}\nabla_a t$. The expansion, vorticity and shear vanish. The acceleration is

$$a_a = \frac{1}{2V} \frac{dV}{dz} \nabla_a z \quad . \quad (3.24)$$

The metric is type D.

First we consider the Killing vector $\xi^a = (\partial/\partial x)^a$. Then we have $\lambda = z$ and thus $F = \cos^2\theta + z^2 \sin^2\theta$. The equation for α_a is $\nabla_{[a}\alpha_{b]} = -\nabla_{[a}y\nabla_{b]}t$ (as well as $\xi^a\alpha_a = 0$). The solution is $\alpha_a = -y\nabla_a t$. It then follows that

$$\eta_a = \nabla_a x - 2 \cos\theta \sin\theta y \nabla_a t \quad . \quad (3.25)$$

The new metric is then given by

$$d\tilde{s}^2 = F \left(-V dt^2 + V dz^2 + z dy^2 \right) + \frac{z}{F} (dx - 2 \cos\theta \sin\theta y dt)^2 \quad . \quad (3.26)$$

Both the seed metric and the new metric have singularities at $z = 0$. However, the new solution does not appear to have any additional singularities. The expansion and the vorticity vanish. The acceleration is

$$\tilde{a}_a = \left(\frac{1}{2V} \frac{dV}{dz} + \frac{z \sin^2 \theta}{F} \right) \nabla_a z \quad . \quad (3.27)$$

The shear has components in the xy and ty directions. Its square is

$$\tilde{\sigma}^{ab} \tilde{\sigma}_{ab} = \frac{2 \cos^2 \theta \sin^2 \theta}{F^3 V} \quad . \quad (3.28)$$

The metric is type I.

We now consider another Killing vector of the Tabensky-Taub metric. Using polar coordinates for the $x - y$ plane we find that the seed metric is

$$ds^2 = -V dt^2 + V dz^2 + z(dr^2 + r^2 d\phi^2) \quad . \quad (3.29)$$

Using the Killing vector $\xi^a = (\partial/\partial\phi)^a$ we find $\lambda = z r^2$ and $\omega = 0$. Therefore $F = \cos^2 \theta + z^2 r^4 \sin^2 \theta$. The equation for α_a is

$$\nabla_{[a} \alpha_{b]} = r \nabla_{[a} r \nabla_{b]} t + 2 V \nabla_{[a} t \nabla_{b]} z \quad (3.30)$$

(as well as $\xi^a \alpha_a = 0$). A solution is

$$\alpha_a = 2 V t \nabla_a z - r t \nabla_a r \quad . \quad (3.31)$$

The vector η_a is then given by

$$\eta_a = \nabla_a \phi + 2 \cos \theta \sin \theta (2 V t \nabla_a z - r t \nabla_a r) \quad . \quad (3.32)$$

The new metric is then given by

$$\begin{aligned} d\tilde{s}^2 = & F \left(-V dt^2 + V dz^2 + z dr^2 \right) \\ & + \frac{z r^2}{F} (d\phi + 2 \cos \theta \sin \theta [2 V t dz - r t dr])^2 \quad . \end{aligned} \quad (3.33)$$

This metric does not have any singularity on the axis (though one must choose the range of ϕ to avoid a conical singularity). The expansion and vorticity vanish. The acceleration is

$$\tilde{a}_a = \left(\frac{1}{2V} \frac{dV}{dz} + \frac{zr^4 \sin^2 \theta}{F} \right) \nabla_a z + \frac{2z^2 r^3 \sin^2 \theta}{F} \nabla_a r \quad . \quad (3.34)$$

The shear has components in the rr , $r\phi$, rz , zz and ϕz directions. Its square is

$$\tilde{\sigma}^{ab} \tilde{\sigma}_{ab} = \frac{2r^2 \cos^2 \theta \sin^2 \theta (r^2 + 4zV)}{F^3 V} \quad . \quad (3.35)$$

The metric is type I.

D. Allnutt seed metric

Our next seed metric is due to Allnutt. [11] It is

$$ds^2 = z^2 \left[-\frac{dt^2}{1+t^2} + f dx^2 + \frac{t^2 (1+t^2)}{f} dy^2 \right] + dz^2 \quad (3.36)$$

where the function f is $f = t^{2\beta} (1+t^2)^{1-\beta}$ and β is a constant. This spacetime is a solution of the Einstein-perfect fluid equations with

$$P = \rho = \frac{\beta (1-\beta)}{8\pi z^2 t^2 (1+t^2)} \quad (3.37)$$

and $u_a = -z(1+t^2)^{-1/2} \nabla_a t$. The vorticity vanishes. The acceleration is $a_a = z^{-1} \nabla_a z$ and the expansion is

$$\Theta = \frac{2t^2 + 1}{zt\sqrt{1+t^2}} \quad . \quad (3.38)$$

The shear has components in the xx , yy and zz directions. Its square is

$$\sigma^{ab} \sigma_{ab} = \frac{2}{3z^2 t^2 (1+t^2)} \left[t^4 + t^2 + 1 + 3\beta(\beta-1) \right] \quad . \quad (3.39)$$

The metric is type D.

Using the Killing vector $\xi^a = (\partial/\partial x)^a$ we find $\lambda = z^2 f$ and thus $F = \cos^2 \theta + z^4 f^2 \sin^2 \theta$.

The equation for α_a is

$$\nabla_{[a} \alpha_{b]} = 2z^2 t \nabla_{[a} t \nabla_{b]} y + 2z (t^2 + \beta) \nabla_{[a} z \nabla_{b]} y \quad (3.40)$$

(as well as $\xi^a \alpha_a = 0$). A solution is $\alpha_a = z^2 (t^2 + \beta) \nabla_a y$. It then follows that

$$\eta_a = \nabla_a x + 2 \cos \theta \sin \theta z^2 (t^2 + \beta) \nabla_a y \quad . \quad (3.41)$$

The new metric is then given by

$$\begin{aligned} d\tilde{s}^2 = F \left[-\frac{z^2}{1+t^2} dt^2 + \frac{z^2 t^2 (1+t^2)}{f} dy^2 + dz^2 \right] \\ + \frac{z^2 f}{F} \left[dx + 2 \cos \theta \sin \theta z^2 (t^2 + \beta) dy \right]^2 \quad . \end{aligned} \quad (3.42)$$

Both the seed metric and the new metric are singular at $z = 0$ and at $t = 0$. However, the new metric does not appear to have any additional singularities. The vorticity vanishes.

The expansion is

$$\tilde{\Theta} = \sqrt{\frac{1+t^2}{F}} \left(\frac{2t^2+1}{zt(1+t^2)} + \frac{z^3 f \dot{f} \sin^2 \theta}{F} \right) \quad (3.43)$$

where an overdot denotes derivative with respect to t . The acceleration is

$$\tilde{a}_a = \left(z^{-1} + 2F^{-1} z^3 f^2 \sin^2 \theta \right) \nabla_a z \quad . \quad (3.44)$$

The shear has components in the xx , yy , zz and xy directions. Its square is

$$\begin{aligned} \tilde{\sigma}^{ab} \tilde{\sigma}_{ab} = \frac{2}{3z^2 t^2 (1+t^2) F} \left[t^4 + t^2 + 1 + 3\beta(\beta-1) \right] \\ + \frac{8}{3} \sin^2 \theta z^2 \frac{f^2}{F^2} \left[\frac{t^2 + \beta}{t} \left(\frac{\dot{F}}{F} + \frac{1-3\beta-t^2}{t(1+t^2)} \right) + \frac{3\cos^2 \theta}{F} \right] \quad . \end{aligned} \quad (3.45)$$

The metric is type I.

E. Kramer seed metric

The final $P = \rho$ seed metric that we shall consider is due to Kramer. [12] It is

$$ds^2 = e^{\alpha^2 r^2} \left(-dt^2 + dr^2 \right) + r^2 d\phi^2 + dz^2 \quad (3.46)$$

where α is a constant. This is a solution of the Einstein-perfect fluid equations with

$$P = \rho = \frac{\alpha^2}{8\pi} e^{-\alpha^2 r^2} \quad . \quad (3.47)$$

The four velocity is $u_a = -e^{\alpha^2 r^2/2} \nabla_a t$. The expansion, shear and vorticity vanish. The acceleration is $a_a = \alpha^2 r \nabla_a r$. The metric is type D. We use the Killing vector $\xi^a = (\partial/\partial\phi)^a$. Then we have $\lambda = r^2$ and thus $F = \cos^2\theta + r^4 \sin^2\theta$. The equation for α_a is $\nabla_{[a}\alpha_{b]} = -2\nabla_{[a}z\nabla_{b]}t$ (as well as $\xi^a\alpha_a = 0$). A solution is $\alpha_a = -2z\nabla_a t$. The new metric is then given by

$$d\tilde{s}^2 = F \left[e^{\alpha^2 r^2} (-dt^2 + dr^2) + dz^2 \right] + \frac{r^2}{F} (d\phi - 4\cos\theta \sin\theta z dt)^2 \quad . \quad (3.48)$$

The new metric has no singularities on the axis, though one must choose the range of ϕ to avoid a conical singularity. The expansion and vorticity vanish. The acceleration is

$$\tilde{a}_a = \left(\alpha^2 r + 2F^{-1}r^3 \sin^2\theta \right) \nabla_a r \quad . \quad (3.49)$$

The shear has components in the $z\phi$ and zt directions. Its square is

$$\tilde{\sigma}^{ab}\tilde{\sigma}_{ab} = \frac{8r^2 e^{-\alpha^2 r^2} \cos^2\theta \sin^2\theta}{F^3} \quad . \quad (3.50)$$

The metric is type I.

IV. $\rho + 3P = 0$ FLUIDS

A. Fluid variables

We now consider the solutions with the equation of state $\rho + 3P = 0$. This equation of state occurs in the Einstein static universe [13], in one of the limiting solutions of the Wahlquist metric [14] due to Vaidya [8,15] and has been suggested as the equation of state for a chaotic array of strings [16]. In generating solutions with the equation of state $\rho + 3P = 0$ we must choose the fluid four-velocity of the seed metric to be parallel to the Killing vector. The fluid four velocity of the new metric will then also be parallel to the Killing vector. This results in considerable simplification of the fluid parameters. The expansion and shear of both the seed solution and the new solution vanish. The density, four-velocity and acceleration transform as follows:

$$\tilde{\rho} = \rho/F \quad , \quad (4.1)$$

$$\tilde{u}^a = \sqrt{F} u^a \quad , \quad (4.2)$$

$$\tilde{a}_a = \left(2F^{-1} \cos^2 \theta - 1 \right) a_a \quad . \quad (4.3)$$

Since we choose seed solutions with zero twist, it follows that the fluid vorticity of the seed solution vanishes. The vorticity of the new metric is

$$\tilde{\omega}_{ab} = 2 \cos \theta \sin \theta \sqrt{-\lambda/F} \nabla_{[a} \alpha_{b]} \quad . \quad (4.4)$$

It then follows that the vorticity vector $\tilde{\Omega}_a := \tilde{\epsilon}_{abcd} \tilde{u}^b \tilde{\omega}^{cd}$ is parallel to the acceleration vector $\tilde{a}_a$.

B. Vaidya seed metric

Our first seed metric is due to Vaidya [15] and is given by

$$ds^2 = -f dt^2 + \frac{dr^2}{(1+cr^2)f} + r^2 (d\psi^2 + \sin^2 \psi d\phi^2) \quad . \quad (4.5)$$

Here

$$f := 1 - \frac{2m}{r} \sqrt{1+cr^2} \quad (4.6)$$

and c and m are constants. This two parameter family of solutions contains the Schwarzschild metric and the Einstein static universe as special cases. The energy density is $\rho = -3cf/(8\pi)$ and the fluid four velocity is $u_a = -\sqrt{f} \nabla_a t$. The acceleration is

$$a_a = \frac{1}{2f} \frac{df}{dr} \nabla_a r \quad . \quad (4.7)$$

The metric is type D. The timelike Killing vector is $\xi^a = (\partial/\partial t)^a$. It then follows that $\lambda = -f$ and therefore that $F = \cos^2 \theta + f^2 \sin^2 \theta$. The equation for α_a is $\nabla_{[a} \alpha_{b]} = -2m \sin \psi \nabla_{[a} \psi \nabla_{b]} \phi$ (as well as $\xi^a \alpha_a = 0$). A solution is $\alpha_a = 2m \cos \psi \nabla_a \phi$. We then find

$$\eta_a = \nabla_a t + 4m \cos \theta \sin \theta \cos \psi \nabla_a \phi \quad . \quad (4.8)$$

The new metric is

$$d\tilde{s}^2 = F \left[\frac{dr^2}{(1+cr^2)f} + r^2 (d\psi^2 + \sin^2\psi d\phi^2) \right] - \frac{f}{F} (dt + 4m \cos\theta \sin\theta \cos\psi d\phi)^2 . \quad (4.9)$$

This metric has a singularity on the axis. The acceleration is

$$\tilde{a}_a = \left(2F^{-1} \cos^2\theta - 1 \right) \frac{1}{2f} \frac{df}{dr} \nabla_a r . \quad (4.10)$$

The nonvanishing components of the vorticity are

$$\tilde{\omega}_{\psi\phi} = -\tilde{\omega}_{\phi\psi} = -2m \cos\theta \sin\theta \sqrt{\frac{f}{F}} \sin\psi . \quad (4.11)$$

After the transformation the pressure and energy density are scaled by F^{-1} from their static values. As in the static case, on the horizon the pressure and scalar curvature are zero. The generated vorticity tensor vanishes on the horizon. The metric is type I.

C. Kramer seed metric

The next seed metric that we will consider is essentially due to Kramer. [12] It is

$$ds^2 = -e^{2x} dt^2 + \frac{1}{e^{2x} \sinh bx} \left[e^{\sqrt{b^2-4}x} d\phi^2 + e^{-\sqrt{b^2-4}x} dz^2 + \frac{dx^2}{\sinh bx} \right] . \quad (4.12)$$

Here b is a constant. The energy density is $\rho = 3b^2 e^{2x}/(32\pi)$ and the fluid four velocity is $u_a = -e^x \nabla_a t$. The acceleration is $a_a = \nabla_a x$. The metric is type I. The Killing vector is $\xi^a = (\partial/\partial t)^a$ and therefore we have $\lambda = -e^{2x}$. It then follows that $F = \cos^2\theta + e^{4x} \sin^2\theta$. The equation for α_a is $\nabla_{[a} \alpha_{b]} = 2 \nabla_{[a} z \nabla_{b]} \phi$ (as well as $\xi^a \alpha_a = 0$). A solution is $\alpha_a = 2z \nabla_a \phi$. We then find

$$\eta_a = \nabla_a t + 4 \cos\theta \sin\theta z \nabla_a \phi . \quad (4.13)$$

The new metric is then given by

$$d\tilde{s}^2 = \frac{F}{e^{2x} \sinh bx} \left[e^{\sqrt{b^2-4}x} d\phi^2 + e^{-\sqrt{b^2-4}x} dz^2 + \frac{dx^2}{\sinh bx} \right] - \frac{e^{2x}}{F} (dt + 4 \cos\theta \sin\theta z d\phi)^2 . \quad (4.14)$$

The solution generating process does not seem to have added any new singularities. The acceleration is

$$\tilde{a}_a = \left(2F^{-1}\cos^2\theta - 1\right) \nabla_a x \quad . \quad (4.15)$$

The nonvanishing components of the vorticity are

$$\tilde{\omega}_{z\phi} = -\tilde{\omega}_{\phi z} = 2\cos\theta\sin\theta e^x F^{-1} \quad . \quad (4.16)$$

The metric is type I.

V. DISCUSSION

Exact solutions to the Einstein field equations are valuable as a tool in the exploration of behavior ranges allowed by the field equations. With the marked increase in numerical calculations, exact solutions are also useful as comparisons for approximate or numerical solutions and as checks for the development of computer codes. In general Einstein's equations for static spaces are far easier to solve than those for stationary spacetimes. One advantage of solution generating methods is that they can be used to find stationary solutions using simple static seed solutions. In this paper, we have used the method of Stephani [7] to generate stationary solutions of the Einstein-perfect fluid field equation from simple static seeds. A possible disadvantage of solution generating is that often extra singularities, especially on an axis, can be introduced. Some of the solutions that we generated have no new singularities and in some cases they appear to be non-singular.

While solution generating techniques have been applied extensively to vacuum spacetimes and to solutions of the Einstein-Maxwell equations [8]; with some notable exceptions [9,10] there is comparatively little work on solution generating involving perfect fluids. The method of reference [10] generates $P = \rho$ perfect fluid solutions from vacuum solutions. It can be applied only to solutions with two spacelike Killing vectors and generates only solutions with zero vorticity. In contrast the Stephani method requires only one Killing vector and is compatible with nonzero vorticity.

There has been recent numerical work on vacuum cosmologies with two spacelike Killing vectors. [17] In this work the known vacuum exact solutions provided useful test cases. The numerical work of reference [17] could easily be generalized to the case of inhomogeneous perfect fluid cosmologies. The exact solutions that we have generated in this paper could be used as test cases for such numerical work.

All of the seed metrics that we have used belong to the Perjes Class II [18]. Since this classification is preserved under the transformation [19] the generated metrics are also members of this class. For the $P = -\rho/3$ solutions, the generated solutions are all shear and expansion free and all have a non-zero vorticity that lies along the static acceleration. This is a characteristic of stationary Perjes II solutions. No assumption about algebraic speciality is made. In the $P = \rho$ case, the fluid parameters of the new solutions are not constrained to be shear free or rigidly rotating. The action of the transformation in this case is on the spacelike Killing vector and all of the generated solutions as well as their seeds have zero vorticity while their shear, expansion and acceleration structure can be quite complex. Solution generating allows access to exact solutions that are difficult to find using other methods. These new solutions illustrate various behaviors allowed by the Einstein-perfect fluid equations and may additionally serve as aids in further numerical work. In some cases they provide insights into the static origins of existing solutions obtained through more arduous methods.

VI. ACKNOWLEDGEMENTS

We thank the University of Michigan for hospitality. D.G. was partially supported by NSF Grant PHY9408439 to Oakland University and by a Cottrell College Science Award of Research Corporation to Oakland University. E.N. Glass was partially supported by an NSERC of Canada grant. Computations were verified using MapleV.3 (Waterloo Maple Software, Waterloo, Ontario) and GRTensorII (P. Musgrave, D.Pollney, and K. Lake, Queens University, Kingston, Ontario).

REFERENCES

- [1] J. Ehlers, *Les theories relativistes de la gravitation* (CNRS, Paris, 1959)
- [2] B. K. Harrison, J. Math. Phys. **9**, 1744 (1968)
- [3] R. Geroch, J. Math. Phys. **12**, 918 (1971)
- [4] D. Kramer and G. Neugebauer, Commun. Math. Phys. **10**, 132 (1968)
- [5] I. Hauser and F. Ernst, J. Math. Phys. **19**, 1316 (1978)
- [6] G. Neugebauer and D. Kramer, Ann. Physik **24** , 62 (1969)
- [7] H. Stephani, J. Math. Phys. **29**, 1650 (1988)
- [8] *Exact Solutions of Einstein's Field Equations* by D. Kramer, H. Stephani, E. Herlt, M. MacCallum and E. Schmutzer, Cambridge University Press (Cambridge, U.K. 1980)
- [9] R. Tabensky and A.H. Taub, Commun. Math. Phys. **29**, 61 (1973)
- [10] J. Wainwright, W.C.W. Ince and B.J. Marshman, Gen. Rel. Grav. **10**, 259 (1979)
- [11] J.A. Allnutt, Gen. Rel. Grav. **13**, 1017 (1981)
- [12] Kramer, Class. Quantum Grav. **5**, 393 (1988)
- [13] *The Large Scale Structure of Space-Time* by S.W. Hawking and G.F.R. Ellis, Cambridge University Press (Cambridge, U.K. 1973)
- [14] H.D. Wahlquist, Phys. Rev. **172**, 1291 (1968)
- [15] P.C. Vaidya, Pramana **8**, 512 (1977)
- [16] A. Vilenkin, Phys. Rev. D **24**, 2082 (1981)
- [17] B. Berger and V. Moncrief, Phys. Rev. D **48**, 4676 (1993); B. Berger, D. Garfinkle and V. Swamy, Gen. Rel. Grav. **27**, 511 (1995)

- [18] Z. Perjés, Int. J. Theor. Phys. **10**, 217 (1974); B. Lukacs, Z. Perjés and A. Sebesstyen, Gen. Rel. Grav. **15**, 511 (1983).
- [19] J.P. Krisch, J. Math. Phys. **29**, 447 (1988).